

Growth Pattern and Condition Factor of *Oreochromis Niloticus* and *Sarotherodon Galilaeus* in Lake Geriyo, Adamawa State

¹H. A., ADEDEJI, ²M. D.HYGINUS, ³T.A. IDOWU, ⁴O. A. SOGBESAN

Department of Fisheries, ModibboAdama University of Technology, Yola

Abstract: The growth pattern of *Oreochromisniloticus* and *Sarotherodongalilaeus* in Lake Geriyo of Adamawa State was investigated. One hundred (100) samples of each species were collected from fishermen catches at the landing site. They were transported to the Department of Fisheries Laboratory for identification, sexing morphometric measurement. The results showed that 58.0% and 56.0 % of the *O.niloticus* and *S.galilaeus* were female respectively while 42.0% and 44.0% were male respectively.. The length-weight relationship of *O.niloticus* showed a strong regression coefficient ($r = 0.944$) ($p < 0.05$) for the combined sex, ($r = 0.903$) ($p < 0.05$) for female and ($r = 0.986$) ($p < 0.05$) for male with b -value of 2.5, 3.1 and 2.8 respectively. The *S.galilaeus* also showed a regression coefficient of $r = 0.970$; $p < 0.05$ for the combined sex, $r = 0.953$; $p < 0.05$ for the female and $r = 0.979$; $p < 0.05$ for the male with b -value of 3.4, 3.1 and 3.6 respectively. *O.niloticus* showed an isometric growth pattern both in male and female. The condition factor in *O.niloticus* are 1.93 and 1.95 for male and female respectively while that of *S.galilaeus* is 1.95 for both sexes. Positive allometric was observed in both species except the male *O.niloticus* that showed negative allometric growth pattern. From the results of this study, the growth pattern of the two species indicated a good healthy environment.

Keywords: Lake Geriyo, Length-weight relationship, *Oreochromisniloticus*, *Sarotherodongalilaeus*.

1. INTRODUCTION

The length-weight relationship of any fish is basically known to measure its growth pattern which is an important component of biological production (Onimisi and Oniye 2010). In fish, size has been known to be more biologically relevant because several ecological and physiological factors are more size-dependent than age-dependent. Also, the length-weight relationships are more important in fisheries biology and fisheries management because they allow and ease the estimation of average weight of fish of a given length (Beyer, 1987). And also it can be used to determine the condition factor in order to express the condition factor of fish in numerical terms (Wootton, 1992). The negative changes in growth rates may result in decreased individual health and reproduction success and increased risk of predation and mortality (Wootton, 1992). When b -value is < 3 , fish is said to have negative allometric growth while > 3 is said to be positive allometric growth (Khaironizan and Norma-Rashid, 2002). Fish is said to be ideal when it maintains its shape as it grows and such fish are said to maintained a three dimensional equality with b -value equal to 3 (isometric growth). Length- weight relationship data can notably provide important clues to climatic and environmental changes, and the change in human subsistence practices (Pauly, 1984; Luff, and Bailey, 2000). Ayoade, (2011), Onimishi and Oniye (2010) and Fafioye and Oluajo (2005) among others have reported on length-weight relationship of different freshwater fish species.

The condition factor (K) which indicates the physiological state of the fish in relation to its welfare (Le Cren, 1951) also provides information when comparing two populations living in certain feeding density, climate and other conditions (Weatherly and Gills 1987). Thus, condition factor is important in understanding the life cycle of fish species, a tool for their adequate management and well-being (Bagenal and Tesch, 1978). It has also been used as an index of growth and feeding intensity (Fagade, 1979); and also influences the reproductive cycle in fish (Welcome, 1979). The works of

Ayoade and Ikulala (2007) and Olurin and Aderibigbe (2006) among others are good literature. This study aims to report the health status of *Oreochromis niloticus* and *Sarotherodon galilaeus* in Lake Gerio, Yola of Adamawa State using their length-weight relationship and condition factors.

2. MATERIALS AND METHODS

The samples were collected from Lake Geriyo, Yola, Adamawa State, Nigeria. It is located on latitude $9^{\circ}81'N$ and longitude $12^{\circ}35'E$ of the equator, the altitude of 185.5m, and the mean annual rainfall is between 900 to 1100mm. The rainy season ranges from 150 – 160 days, usually from May to October. The average annual temperature of the area is $26.8^{\circ}C$ with average annual rainfall of 458mm (AADIL, 2011).

The fish samples were identified using Field guide to Nigeria Freshwater Fishes by Olaosebikan and Raji (2013). The collection was done weekly for a period of two months from the fishermen at the landing sites of Lake Geriyo and immediately transported to the laboratory in an ice box.

The total and standard lengths were measured using a measuring board of 50cm range and all measurements were determined to the nearest centimetres. Body weight was measured using an Ohaus digital weighing balance (Mettler Instrument). Fish samples were differentiated into separate sexes using gonadal differentiation. The numbers of specimens in each sex were recorded.

Estimation of species length-weight relationship was done using the formula $W=aL^b$ (Ricker, 1975), which is transformed into natural logarithmic form $\log W = \log a + b \log L$, where W is the total weight of fish (grams), L is the total length of fish (centimeters), a is constant (intercept) and b is the regression coefficient (slope) (Bagenal and Tesch, 1978). While condition factor (K) was computed using the formula: $K=100W/L^3$ (Pauly, 1983). Means of data generated were determined using descriptive statistics while significant levels of means were compared at $p<0.05$ using One-way ANOVA. Means predictions were done using Regression equation.

3. RESULTS AND DISCUSSION

Table 1 showed the mean total length, mean total weight, length-weight relationship parameters and condition factor of *O. niloticus* and *S. galilaeus*. The mean total length recorded from the study was 15.23 ± 1.6 cm (combined sex), 14.74 ± 1.71 (male), 15.58 ± 1.51 (female) and 16.23 ± 1.41 (combined), 15.72 ± 1.51 (male), 16.63 ± 1.22 (female) for *O. niloticus* and *S. galilaeus* respectively. The mean body weight was 65.97 ± 20.87 g (combined), 62.49 ± 20.30 (male), 68.49 ± 21.29 (female) and 85.28 ± 23.90 g (combined), 78.00 ± 26.28 (male), 90.99 ± 20.55 (female) for *O. niloticus* and *S. galilaeus* respectively. The length-weight relationship of *O. niloticus* in Lake Geriyo showed strong correlation ($r=0.986$) for male, ($r=0.903$) for female and ($r=0.9044$) for the combined sex and likewise *S. galilaeus* ($r=0.979$) for male, ($r=0.953$) for female and ($r=0.970$) for combined sex indicating an increase in weight as the length increases. The male *O. niloticus* showed negative allometric growth ($b=2.8$) while the female showed positive allometric growth ($b=3.1$). The *S. galilaeus* species showed positive allometric growth both in male ($b=3.5$) and female ($b=3.2$). The mean condition factor (K) in this study showed k -value higher than 1 which is an indication of good feeding and favorable environmental conditions as condition factor is known to describe the well-being of a species or population of species in an environment (Ikomi and Odum, 1998).

Table 1: Mean Total Length, Mean Total Weight, Length-Weight relationship parameters and Condition factor of *O. niloticus* and *S. galilaeus*

	<i>Oreochromis niloticus</i>			<i>Sarotherodon galilaeus</i>		
	Male	Female	Combined	Male	Female	Combined
BW(g)	62.49 ± 20.30^c	68.49 ± 21.29^c	65.97 ± 20.87^c	78.00 ± 26.28^c	90.99 ± 20.55^a	85.28 ± 23.90^a
TL(cm)	14.74 ± 1.71^a	15.58 ± 1.51^a	15.23 ± 1.6^a	15.72 ± 1.51^a	16.63 ± 1.22^a	16.23 ± 1.41^a
SL(cm)	11.51 ± 1.38^a	12.03 ± 1.28^a	11.81 ± 1.32^a	12.11 ± 1.43^a	12.71 ± 0.99^a	12.4 ± 1.22^a
b	2.8	3.1	2.5	3.5	3.2	3.4
r	0.986	0.903	0.944	0.979	0.953	0.970
K	1.93	1.95	1.94	1.95	1.95	1.93
n	42	58	100	44	56	100

Keys: BW – Body weight; TL – Total length; SL – Standard length

REFERENCES

- [1] Ayoade, A. A. (2011). The length-weight relationship of *Schilbemystus* (Linne,1766) from two man-made lakes in south-western Nigeria. *The Zoologist*, 9:38-43
- [2] Ayoade, A. A. and Ikulala, A. O. O. (2007).Length weight relationship, condition factor and stomach contents of *Hemichromisbimaculatus*, *Sarotherodonmelanotheron* and *Chromidotilapiaquentheri* (Perciformes: Cichlidae) in Eleiyele Lake, Southwestern Nigeria. *Rev. Biol. Trop. (Int. Journal of Tropical Biology)* . Vol. 55 (3-4): 969-977
- [3] Bagenal, T.B. and A.T. Tesch., (1978).Conditions and growth patterns in fresh water habitats.*Blackwell Scientific Publications, Oxford*. Pp. 75-89
- [4] Beyer, J.E., (1987).On length-weight relationship computing the mean weight of the fish of a given length class.*fishbyte*, 5(1): 11-13.
- [5] Fafioye,O.O. and Oluajo,O.A. (2005) length-weight relationships of five fish species in Epe Lagoon, Nigeria. *African Journal Biotechnology*, 4 (7):749-751.
- [6] Fagade, S.O. (1979). Observation of the biology of two species of *Tilapia* from the Lagos lagoon Nigeria.*Bull. Inst. Fond Afr. Nore., (series. A) 41: 627-658*.
- [7] Ikomi, R. B. and Odum, O. (1998).Studies on aspect of the ecology of catfish *Chrysichthysauratus* Geoffrey St. Hilaire (Osteichthys; Bagridae) in the river Benin (Niger Delta, Nigeria).*FisheriesResearch*, 35:209-218
- [8] Khaironizam, M. Z. and Norman-Rashid, Y. (2002). Length-weight relationship of mudskippers (Gobiidae: Oxudercinae) in the coastal areas of Selangor, Malaysia. International Centre for Living Aquatic Resources Management.*World Fish Centre Quarterly*, 25:20-22
- [9] Le cren, E.D., (1951). The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Percafluviatilis*). *J. Anim. Ecol.*, 20(2): 201-219.
- [10] Luff, R. M and Bailey, G. N. (2000). Analysis of size changes and incremental growth structures in African catfish *Synodontisschall* (schall) from Tell el-Amarna, Middle Egypt. *Journal Archeology Sciences*, 27: 821-835.
- [11] Olaosebikan and Raji (2013).Field guide to Nigeria Freshwater Fishes.
- [12] Olurin, K.B. and Aderigbe, O.A. (2006). Length-weight relationship and condition factor of pond-reared juveniles *Oreochromisniloticus* .*World J.Zoology* , 1(2):82-85.
- [13] Onimishe, H. U. and Oniye, S. J. (2010). Length-weight relationship and condition factor of *Auchenoglanisoccidentalis* in Zaria reservoir, Nigeria. *The Zoologist*, Vol. 8:52-56
- [14] Pauly, D and Gayanilo Jr. F. C. (1997). A Bee: an alternative approach to estimating the parameters of length-weight relationship from length-frequency samples and their bulk weights. *International centre for Living Aquatic Resources Management Quarterly*, pp.168
- [15] Pauly, D. (1984). Fish population dynamic in tropical waters: a manual for use with programmable calculators. Naga, ICLARM Qtr., 5-95.
- [16] Ricker W.E. (1975). Computation and interpretation of biological statistics of fish population, *Bulletin of the Fisheries Research Board of Canada* 191: 1-382.
- [17] Weatherly,A.H. and Gill,H.S. (1987): *The biology of fish growth* ,AcademicPress, London, 433-443.
- [18] Welcome, R. L.(1979). Fisheries Ecology of flood plain rivers.*London long man press*.pp: 317.